

Asthma Treatment Plan

(This asthma action plan meets NJ Law N.J.S.A. 18A:40-12.8) (Physician's Orders)



"Your Pathway to Asthma Control"
Original PACNJ approved Plan available at
www.pacnj.org



(Please Print)

Name	Date of Birth	Effective Date
Doctor	Parent/Guardian (if applicable)	Emergency Contact
Phone	Phone	Phone

HEALTHY



You have all of these:

- Breathing is good
- No cough or wheeze
- Sleep through the night
- Can work, exercise, and play

And/or Peak flow above _____

Take daily medicine(s). All metered dose inhalers (MDI) to be used with spacers.

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
<input type="checkbox"/> Advair® 100, 250, 5001 inhalation twice a day
<input type="checkbox"/> Advair® HFA 45, 115, 2302 puffs MDI twice a day
<input type="checkbox"/> Asmanex® Twisthaler® 110, 2201 - 2 inhalations a day
<input type="checkbox"/> Flovent® 44, 110, 2202 inhalations twice a day
<input type="checkbox"/> Flovent® Diskus® 50 mcg1 inhalation twice a day
<input type="checkbox"/> Pulmicort Flexhaler® 90, 1801 - 2 inhalations once or twice a day
<input type="checkbox"/> Pulmicort Respules® 0.25, 0.5, 1.01 unit nebulized once or twice a day
<input type="checkbox"/> Qvar® 40, 802 inhalations twice a day
<input type="checkbox"/> Singulair 4, 5, 10 mg1 tablet daily
<input type="checkbox"/> Symbicort® 80, 1602 puffs MDI twice a day
<input type="checkbox"/> Other	

Remember to rinse your mouth after taking inhaled medicine.

If exercise triggers your asthma, take this medicine _____ minutes before exercise.

CAUTION



You have any of these:

- Exposure to known trigger
- Cough
- Mild wheeze
- Tight chest
- Coughing at night
- Other: _____

And/or Peak flow from _____ to _____

Continue daily medicine(s) and add fast-acting medicine(s).

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
<input type="checkbox"/> Accuneb® 0.63, 1.25 mg1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Albuterol 1.25, 2.5 mg1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Albuterol <input type="checkbox"/> Pro-Air <input type="checkbox"/> Proventil® .2	puffs MDI every 4 hours as needed
<input type="checkbox"/> Ventolin® <input type="checkbox"/> Maxair <input type="checkbox"/> Xopenex® .2	puffs MDI every 4 hours as needed
<input type="checkbox"/> Xopenex® 0.31, 0.63, 1.25 mg1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Increase the dose of, or add:	

➡ If fast-acting medicine is needed more than 2 times a week, except before exercise, then call your doctor.

EMERGENCY



Your asthma is getting worse fast:

- Fast-acting medicine did not help within 15-20 minutes
- Breathing is hard and fast
- Nose opens wide
- Ribs show
- Trouble walking and talking
- Lips blue • Fingernails blue

And/or Peak flow below _____

Take these medicines NOW and call 911. Asthma can be a life-threatening illness. Do not wait!

<input type="checkbox"/> Accuneb® 0.63, 1.25 mg1 unit nebulized every 20 minutes
<input type="checkbox"/> Albuterol 1.25, 2.5 mg1 unit nebulized every 20 minutes
<input type="checkbox"/> Albuterol <input type="checkbox"/> Pro-Air <input type="checkbox"/> Proventil® .2	puffs MDI every 20 minutes
<input type="checkbox"/> Ventolin® <input type="checkbox"/> Maxair <input type="checkbox"/> Xopenex® .2	puffs MDI every 20 minutes
<input type="checkbox"/> Xopenex® 0.31, 0.63, 1.25 mg1 unit nebulized every 20 minutes
<input type="checkbox"/> Other	

Triggers

Check all items that trigger patient's asthma:

- Chalk dust
- Cigarette Smoke & second hand smoke
- Colds/Flu
- Dust mites, dust, stuffed animals, carpet
- Exercise
- Mold
- Ozone alert days
- Pests - rodents & cockroaches
- Pets - animal dander
- Plants, flowers, cut grass, pollen
- Strong odors, perfumes, cleaning products, scented products
- Sudden temperature change
- Wood Smoke
- Foods:

Other: _____

This asthma treatment plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs.

The Pediatric/Adult Asthma Coalition of New Jersey, sponsored by the American Lung Association of New Jersey, and this publication are supported by a grant from the New Jersey Department of Health and Senior Services (NJDHSS), with funds provided by the U.S. Centers for Disease Control and Prevention (USCDC) under Cooperative Agreement #U59CE00026-2. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NJDHSS or the USCDC.

Although this document has been funded wholly or in part by the United States Environmental Protection Agency under Agreements #68028401-4 and #68025002-1 to the American Lung Association of New Jersey, it has not gone through the Agency's publication review process and therefore, may not necessarily reflect the views of the Agency and no official endorsement should be inferred.

EFFECTIVE MARCH 2008

Permission to reproduce blank form. Approved by the New Jersey Thoracic Society

FOR MINORS ONLY:

- This student is capable and has been instructed in the proper method of self-administering of the inhaled medications named above in accordance with NJ Law.
- This student is not approved to self-medicate.

PHYSICIAN/APN/PA SIGNATURE _____ DATE _____

PARENT/GUARDIAN SIGNATURE _____

PHYSICIAN STAMP

Make a copy for patient and for physician file. For children under 18, send original to school nurse or child care provider.